

A simple $O(mn)$ algorithm for recognizing Hamming graphs

W. Imrich
Institut für Mathematik und Angewandte Geometrie
Montanuniversität Leoben
A-8700 Leoben
Austria

S. Klavžar *
University of Maribor
PF, Koroška cesta 160
62000 Maribor
Slovenia

Abstract

We show that any isometric irredundant embedding of a graph into a product of complete graphs is the canonical isometric embedding. This result is used to design a simple $O(mn)$ algorithm for recognizing Hamming graphs.

1 Introduction

All graphs considered in this paper are finite undirected graphs without loops and multiple edges. Throughout the paper, for a given graph G , let n and m stand for the number of its vertices and edges, respectively.

Graphs that can be embedded isometrically in a Cartesian product of complete graphs are called Hamming graphs. Interest in Hamming graphs has been decisively stimulated by the work of Graham and Pollak [10, 11]

*This work was supported in part by the Ministry of Science and Technology of Slovenia under the grant P1-0206-101-92.

in communication theory and Firsov [8] in linguistics. In biology, Hamming graphs appear as “quasi-species” [6].

Several algorithms for recognizing Hamming graphs have been proposed. The first algorithm is due to Winkler [16]. Its running time is bounded by $O(n^5)$. Aurenhammer, Formann, Idury, Schäffer and Wagner [2] improved Winkler’s algorithm to run in $O(D(m, n) + n^2)$ time, where $D(m, n)$ is the time needed to compute the distance matrix of the graph. Wilkeit [14] presented another algorithm running in $O(n^3)$ time.

In 1985 Graham and Winkler [12] published a fundamental paper on isometric embeddings into Cartesian product graphs. Using their results together with a result of Aurenhammer and Hagauer [4] one can give an $O(mn)$ algorithm to recognize Hamming graphs. Aurenhammer and Hagauer [3] also proposed another algorithm of the same complexity to recognize binary Hamming graphs.

In this paper we present an $O(mn)$ algorithm to recognize Hamming graphs. Our algorithm is also based on the theory of isometric embeddings into Cartesian product graphs. However, our approach needs a minimum of theory and the algorithm itself is simple and straightforward. We have taken care that everything up to and including Section 4 is self-contained. In particular, this includes all results about binary Hamming graphs.

In the next section we state the necessary definitions and recall a connection between Hamming graphs and the Cartesian product of graphs. In Section 3 we introduce the relation Θ and reprove a characterization of binary Hamming graphs. Section 4 follows with an algorithm for the recognition of binary Hamming graphs. In the last section we prove that any isometric irredundant embedding of a graph into a product of complete graphs is the canonical isometric embedding. With the aid of this result we extend the algorithm for binary Hamming graphs to all Hamming graphs.

2 Hamming graphs and the Cartesian product

Let Σ be a finite alphabet and let w_1 and w_2 be words of equal length over Σ . Then the *Hamming distance* between w_1 and w_2 , $H(w_1, w_2)$, is the number of positions k in w_1 and w_2 such that the k -th symbol in w_1 differs from the k -th symbol in w_2 . A graph G is called a *Hamming graph*, if each vertex $v \in V(G)$ can be labelled by a word of fixed length, $a(v)$, such that $H(a(u), a(v)) = d_G(u, v)$ for all $u, v \in V(G)$. Here $d_G(u, v)$ denotes the usual shortest path distance in G between u and v . In particular, if $\Sigma = \{0, 1\}$, we call G a *binary Hamming graph*.

The *Cartesian product* $G \square H$ of graphs G and H is the graph with vertex set $V(G) \times V(H)$ and $(a, x)(b, y) \in E(G \square H)$ whenever $ab \in E(G)$ and $x = y$, or $a = b$ and $xy \in E(H)$. The Cartesian product is commutative, associative and K_1 is a unit. Also, $G \square H$ is connected if and only if both G and H are connected. For $G_1 \square G_2 \square \cdots \square G_k$ we shall also write $\prod_{i=1}^k G_i$.

For $G = \prod_{i=1}^k G_i$ let $p_i : V(G) \rightarrow V(G_i)$, $i \in \{1, 2, \dots, k\}$, be the natural projection of G onto the i -th factor G_i , i.e. for $v = (v_1, v_2, \dots, v_k) \in V(G)$ we set $p_i(v) = v_i \in V(G_i)$. For $X \subseteq V(G)$ let $p_i(X) = \{p_i(x) \mid x \in X\}$. In particular, for $e = uv \in E(G)$ let $p_i(e) = \{p_i(u), p_i(v)\}$. We also introduce a *product coloring* $c : E(G) \rightarrow \{1, 2, \dots, k\}$ (*with respect to the product representation*) as follows. For $uv \in E(G)$ we set $c(uv) = i$ if and only if u and v differ in coordinate i . Clearly c is a mapping from $E(G)$ into $\{1, 2, \dots, k\}$. It is not an edge coloring in the usual sense, because incident edges may have the same color.

A subgraph H of a graph G is an *isometric* subgraph, if $d_H(u, v) = d_G(u, v)$ for all $u, v \in V(H)$. In addition, if $\alpha : V(H) \rightarrow V(G)$ maps edges into edges and $\alpha(H)$ is an isometric subgraph of G , we call α an *isometric embedding* of H into G .

The following observation is a starting point in studying isometric sub-

graphs of Cartesian products of graphs.

Lemma 2.1 *For $G = \prod_{i=1}^k G_i$, $k \geq 1$, let $u = (u_1, u_2, \dots, u_k)$ and $v = (v_1, v_2, \dots, v_k)$ be arbitrary vertices of G . Then $d_G(u, v) = \sum_{i=1}^k d_{G_i}(u_i, v_i)$.*

With Lemma 2.1 it is an easy exercise to characterize Hamming graphs:

Theorem 2.2 *A graph G is a Hamming graph if and only if G is an isometric subgraph of a Cartesian product of complete graphs. \square*

Theorem 2.2 in particular implies that binary Hamming graphs are isometric subgraphs of hypercubes (the k -dimensional hypercube Q_k , $k \geq 1$, is the Cartesian product of k copies of K_2).

3 The relation Θ

In this section we present a characterization of binary Hamming graphs which is essential for a fast recognition algorithm.

Let G be a connected graph. Define a relation Θ on $E(G)$ as follows. If $e = uu' \in E(G)$ and $e' = vv' \in E(G)$, then $e\Theta e'$ if and only if

$$d(u, v) + d(u', v') \neq d(u, v') + d(u', v).$$

This relation, which was first introduced in an alternative form in [5], plays a central role in our investigation. The relation Θ is well-defined, reflexive and symmetric, yet it need not be transitive. We denote its transitive closure by Θ^* .

Lemma 3.1 *Let P be a shortest path in a graph G . Then no two different edges of P are in relation Θ .*

Proof. Let $u_0u_1 \cdots u_m$ be a shortest path, and let $e = u_iu_{i+1}$, $e' = u_ju_{j+1}$, $i < j$. Then

$$d(u_i, u_j) + d(u_{i+1}, u_{j+1}) = (d(u_{i+1}, u_j) + 1) + (d(u_i, u_{j+1}) - 1),$$

hence e is not in the relation Θ with e' . □

It follows from Lemma 3.1 that for a tree on n vertices Θ^* consists of $n - 1$ equivalence classes, each containing a single edge. Lemma 3.1 also implies that two adjacent edges are in relation Θ if and only if they lie in a common triangle.

Lemma 3.2 *Suppose P is a path connecting the endpoints of an edge e . Then P contains an edge f with $e\Theta f$.*

Proof. Let $u_0u_1 \cdots u_mu_0$ be a closed path with $e = u_mu_0$ and $e_i = u_{i-1}u_i$ for $i = 1, 2, \dots, m$. Set

$$\mu(e, e_i) = d(u_m, u_{i-1}) + d(u_0, u_i) - d(u_m, u_i) - d(u_0, u_{i-1})$$

and consider

$$s = \sum_{i=1}^m \mu(e, e_i).$$

Clearly $s = 2$, which means that at least one of the summands $\mu(e, e_i) \neq 0$, i.e. $e\Theta e_i$. □

For an edge uv of a graph G let

$$V_{uv} = \{ w \mid w \in V(G), d_G(w, u) < d_G(w, v) \}.$$

Lemma 3.3 *Let $e = uv$ be an edge of a connected bipartite graph G and let*

$$E_e = \{ f \mid f \in E(G), e\Theta f \}.$$

Then $G \setminus E_e$ has exactly two components. Furthermore, they are induced by the vertex sets V_{uv} and V_{vu} .

Proof. $G \setminus E_e$ is disconnected by Lemma 3.2.

Let $w \in V_{uv}$ and let P be a shortest $u - w$ path. Then vP is a shortest $v - w$ path. Hence, using Lemma 3.2 again, no edge of P is in relation Θ with e . It follows that P connects u and w in $G \setminus E_e$.

Since G is bipartite, $d(w, u) \neq d(w, v)$. Therefore V_{uv} and V_{vu} form a partition of $V(G)$. In addition, no edge $ww' \in E(G)$, where $w, w' \in V_{uv}$, is in relation Θ with e , and the proof is complete. \square

For Cartesian product graphs Lemma 2.1 immediately implies:

Lemma 3.4 *Let $G = \prod_{i=1}^k G_i$ and let $e, e' \in E(G)$.*

(i) If $c(e) = c(e') = i$ and $p_i(e) = p_i(e')$ then $e\Theta e'$.

(ii) If $c(e) \neq c(e')$ then $e\Theta e'$ does not hold.

Consider the product $G \square H$ of two graphs. Then Lemma 3.4 (i) claims that if $ab \in E(G)$, then $(a, x)(b, x)$ is in relation Θ with $(a, y)(b, y)$, for any $x, y \in V(H)$. Lemma 3.4 (ii) on the other hand says that if $ab \in E(G)$ and $xy \in E(H)$, then $(a, z)(b, z)$ is not in relation Θ with $(c, x)(c, y)$, for any $c \in V(G)$ and $z \in V(H)$, i.e. edges with different colors with respect to the product coloring are not in relation Θ . In other words, every color class (with respect to the product coloring) is the union of one or more equivalence classes with respect to Θ^* .

Theorem 3.5 [16, Winkler] *A graph G is a binary Hamming graph if and only if G is bipartite and $\Theta^* = \Theta$.*

Proof. Assume G is an isometric subgraph of a hypercube. Then G is clearly bipartite.

Define a relation R on $E(G)$ as follows. For $e, e' \in E(G)$ let eRe' if and only if $c(e) = c(e')$. Let $c(e) = c(e') = i$. Then $p_i(e) = p_i(e') = \{0, 1\}$ and hence, by Lemma 3.4 (i), $e\Theta e'$. Furthermore, by Lemma 3.4, e and e' are

not related by Θ if $c(e) \neq c(e')$. It follows $R = \Theta$. As R is transitive, we conclude $\Theta^* = \Theta$.

Conversely, let G be bipartite and let $\Theta^* = \Theta$. Let $e_1 = x_1y_1, e_2 = x_2y_2, \dots, e_k = x_ky_k, k \geq 1$ be representatives of each equivalence class of Θ^* . Define an embedding $\alpha : V(G) \rightarrow Q_k = \{0, 1\}^k$ in the following way. Let $v \in V(G)$. For $i = 1, 2, \dots, k$ let the i -th coordinate of $\alpha(v)$ be 0 if $v \in V_{x_iy_i}$ and 1 if $v \in V_{y_ix_i}$. We claim that α is an isometric embedding.

Let $uv \in E(G)$ and assume uv belongs to the equivalence class of e_i . By Lemma 3.3, $\alpha(u)$ and $\alpha(v)$ differ in the i -th coordinate. Furthermore, if $j \neq i$ then $d(u, x_j) + d(v, y_j) = d(u, y_j) + d(v, x_j)$. Thus $\alpha(u)$ and $\alpha(v)$ have same j -th coordinate. It follows that α maps edges to edges.

As G is bipartite and $\Theta = \Theta^*$, no two adjacent edges are in the same equivalence class. Furthermore, if P is a shortest path between u and v then, by Lemma 3.1, $\alpha(u)$ and $\alpha(v)$ differ in just as many coordinates as the number of edges of P , i.e., the distance $d_G(u, v)$. \square

4 Recognizing binary Hamming graphs

In a direct implementation of Theorem 3.5 we must check for all pairs of edges whether they are in relation Θ . For a given graph G this leads to an algorithm with time complexity $O(m^2)$. In order to improve this complexity, we introduce relation Θ_1 , which is due to Feder [7].

Let T be a spanning tree of a graph G . Then edges $e, e' \in E(G)$ are in the relation Θ_1 if and only if they are in the relation Θ and if at least one of the edges e and e' belongs to T . Let E_1, E_2, \dots, E_k be the equivalence classes of the relation Θ_1^* . For $i = 1, 2, \dots, k$ let G_i denote the graph $(V(G), E(G) \setminus E_i)$ and let $C_{i,1}, C_{i,2}, \dots, C_{i,m_i}$ denote the connected components of G_i . Form the graphs $G_i^*, i = 1, 2, \dots, k$, by letting $V(G_i^*) = \{C_{i,1}, C_{i,2}, \dots, C_{i,m_i}\}$ and taking $C_{i,j}C_{i,j'}$ to be an edge of G_i^* if

some edge in E_i joins a vertex in $C_{i,j}$ to a vertex in $C_{i,j'}$.

We now define the natural contraction $\alpha_i : V(G) \rightarrow V(G_i^*)$ by setting $\alpha_i(v) = C_{i,j}$ if $v \in C_{i,j}$. Finally, we obtain a mapping

$$\alpha : V(G) \rightarrow \prod_{i=1}^k G_i^* \quad (*)$$

by setting

$$\alpha(v) = (\alpha_1(v), \alpha_2(v), \dots, \alpha_k(v)). \quad (*)$$

It is important for our algorithm that $k \leq n-1$. Indeed, let $e = uv \in E_i$. By Lemma 3.2 any path in G from u to v must traverse at least one edge from E_i . This is in particular true for the path between u and v in T , hence there is at least one edge of T belonging to E_i .

This also means that $\Theta = \Theta_1^*$ in a binary Hamming graph. For, suppose $e\Theta f$. Then there is an $e' \in T$ with $e'\Theta e$. Since $\Theta = \Theta^*$ we also have $e'\Theta f$ and $e\Theta_1^* f$. Hence $\Theta \subseteq \Theta_1^*$. Since $\Theta_1 \subseteq \Theta$ we also have $\Theta_1^* \subseteq \Theta^* = \Theta$, which proves the assertion.

Algorithm BHG

Input: a connected graph G .

Output: TRUE, and a labeling, if G is a binary Hamming graph. FALSE, otherwise.

1. If G is not bipartite then return FALSE and stop.
2. Compute Θ_1^* .
3. Compute G_i , $i = 1, 2, \dots, k$ and $\alpha(v)$, $v \in V(G)$.
4. For $i = 1, 2, \dots, k$, compute m_i , i.e., the number of components in G_i . If for some i , $m_i > 2$, then return FALSE and stop.
5. Return TRUE and the labeling of G obtained in step 3.

Theorem 4.1 *Algorithm BHG correctly recognizes binary Hamming graphs and can be implemented to run in $O(mn)$ time using $O(m)$ space.*

Proof. Suppose that G is a binary Hamming graph. Then $\Theta = \Theta_1^*$ and every G_i has exactly two components by Lemma 3.3. It thus remains to show that α is an isomorphism, if every G_i has exactly two components.

Let P be a shortest path between u and v in G . By Lemma 3.1 no two edges of P are in the relation Θ and thus all edges of P belong to different Θ_1^* classes. This means that the labels of u and v differ in just as many coordinates as the number of edges of P , i.e. the distance $d_G(u, v)$. This proves the correctness of the algorithm.

Concerning the running time we first note that it is trivial to check bipartiteness in $O(m)$ time and space.

Let T be a spanning tree of G and let uv be an edge of T . Then we can calculate the distances from u and v to all other vertices in $O(m)$ time and $O(m)$ space. Hence we get all the edges in G related to e under Θ_1 in the same time, and in time $O(mn)$ over all edges in T . To get the equivalence classes of Θ_1^* we merge equivalence classes of two edges whenever we determine that they are in the relation Θ_1 . During the execution of the procedure there will be at most $m - 1$ such union operations and $m(n - 1)$ find operations. It is well-known that these can be done in $O(nm)$ time using $O(m)$ space (see, for example, [1]).

When E_i is known, it is easy to construct the graph G_i in $O(m)$ time. As $k \leq n - 1$, all the G_i can be obtained in $O(mn)$ time. Since every edge in G_i corresponds to an edge of G , we get all the G_i using $O(m)$ space. In addition, we don't need to store the complete information on $\alpha(v)$, it is enough to store the value $\alpha_i(v)$ if it is different from $\alpha_i(u)$ for some $uv \in E(G)$. \square

Graham [9] proved that there exists a fixed small c such that $m \leq cn \log n$ for any subgraph of a hypercube. Thus:

Corollary 4.2 *Algorithm BHG can be implemented to run in $O(n^2 \log n)$ time using $O(n \log n)$ space.*

5 Recognizing Hamming graphs

We are going to modify Algorithm BHG to recognize general Hamming graphs.

Let α be the mapping $(*)$ defined in Section 4. Call an isometric embedding $\beta : G \rightarrow \prod_{i=1}^m H_i^*$ *irredundant* if $|H_i| \geq 2$, $i = 1, 2, \dots, m$, and for all $h \in V(H_i)$, h occurs as a coordinate value of the image of some $g \in V(G)$. This is the principal result in the theory of isometric embeddings into Cartesian product graphs:

Theorem 5.1 [12, Graham and Winkler] *The mapping α is an isometric embedding of G into $\prod_{i=1}^k G_i^*$, the so-called canonical embedding. Furthermore, the embedding α is irredundant and has the largest possible number of factors among all irredundant isometric embeddings of G .*

Graham and Winkler [12] proved Theorem 5.1 for the relation Θ , while Feder [7] showed $\Theta_1^* = \Theta^*$.

For the isometric embeddings into products of complete graphs we have:

Theorem 5.2 [16, Winkler] *Any two isometric embeddings of a graph into products of complete graphs are equivalent.*

Equivalence in Theorem 5.2 essentially means that equivalent isometric embeddings can be obtained from one another by discarding unused factors, permuting factors, and permuting vertices within a factor.

The following theorem is crucial for our algorithm.

Theorem 5.3 *Let $\beta : G \rightarrow \prod_{i=1}^m H_i$ be an isometric irredundant embedding of a graph G into a product of complete graphs H_i . Then this embedding is the canonical isometric embedding.*

Proof. By Theorem 5.2 we have to show that any two edges e, e' in $\beta(G)$ are in the relation Θ^* if they have the same color with respect to the product coloring of $\prod_{i=1}^m H_i$.

Consider now i -layers U and V with respect to H_i . We claim that $p_i(\beta(G) \cap U) \subseteq p_i(\beta(G) \cap V)$ or vice versa. If this is not the case, there are vertices $u, u' \in U$ and $v, v' \in V$ such that

$$u \in \beta(G) \cap U, u' \notin \beta(G) \cap U$$

and

$$v \in \beta(G) \cap V, v' \notin \beta(G) \cap V,$$

where $p_i(u) = p_i(v')$, $p_i(u') = p_i(v)$ (see Fig. 1).




Figure 1: Layers U and V

Suppose the distance between U and V in $\prod_{i=1}^m H_i$ is k . Then $k + 1 = d_H(u, v) = d_{\beta(G)}(u, v)$, which is only possible if $u' \in \beta(G)$ or $v' \in \beta(G)$. This proves the claim.

Thus, let the i -colored edges e, e' in $\beta(G)$ be in the i -layers U and V , respectively. If $p_i(\beta(G) \cap U) \subseteq p_i(\beta(G) \cap V)$, then by Lemma 3.4 (i) there is an edge e'' in $\beta(G) \cap V$ with $e\Theta e''$. Since $\beta(G) \cap V$ is complete we have $e''\Theta e'$, and hence $e\Theta^* e'$. \square

Algorithm HG

Input: a connected graph G .

Output: TRUE, and a labeling, if G is a Hamming graph. FALSE, otherwise.

1. Compute Θ_1^* .
2. Compute G_i , $i = 1, 2, \dots, k$ and $\alpha(v)$, $v \in V(G)$.
3. If for some $i = 1, 2, \dots, k$, G_i^* is not a complete graph, then return FALSE and stop.
4. Return TRUE and the labeling of G obtained in step 2.

Theorem 5.4 *Algorithm HG correctly recognizes Hamming graphs and can be implemented to run in $O(mn)$ time using $O(m)$ space.*

Proof. Correctness of the algorithm follows from Theorem 5.3.

The complexity can be argued as in the proof of Theorem 4.1 with the exception of Step 3. Every edge of a factor graph G_i^* correspond to an edge of G and furthermore, this correspondence is injective. It follows $\sum_{i=1}^k |E(G_i^*)| \leq m$. Hence to implement Step 3 in the desired time and space it is sufficient to count the number of edges in the G_i^* 's. \square

References

- [1] A. Aho, J. Hopcroft and J. Ullman, The Design and Analysis of Computer Algorithms (Addison-Wesley, Reading, 1974).
- [2] F. Aurenhammer, M. Formann, R. Idury, A. Schäffer and F. Wagner, Faster isometric embeddings in products of complete graphs, Discrete Appl. Math., to appear.
- [3] F. Aurenhammer and J. Hagauer, Recognizing binary Hamming graphs in $O(n^2 \log n)$ time, in: Proc. 16th Int. Workshop on Graph Theoretical Concepts in Computer Science, Lecture Notes in Comput. Sci., Vol. 484 (Springer, New York, 1991) 90–98.
- [4] F. Aurenhammer and J. Hagauer, Computing equivalence classes among the edges of a graph with applications, Discrete Math. 109 (1992) 3–12.

- [5] D. Djoković, Distance preserving subgraphs of hypercubes, *J. Combin. Theory Ser. B* 14 (1973) 263–267.
- [6] M. Eigen and R. Winkler-Oswattisch, Transfer-DNA: the early adaptor, *Naturwissenschaften* 68 (1981) 217–228.
- [7] T. Feder, Product graph representations, *J. Graph Theory* 16 (1992) 467–488.
- [8] V. Firsov, Isometric embedding of a graph in a Boolean cube (Russian), *Kibernetika* 1 (1965) 95–96.
- [9] R. Graham, On primitive graphs and optimal vertex assignments, *Ann. New York Acad. Sci.* 175 (1970) 170–186.
- [10] R. Graham and H. Pollak, On the addressing problem for loop switching, *Bell System Tech. J.* 50 (1971) 2495–2519.
- [11] R. Graham and H. Pollak, On embedding graphs in squashed cubes, in: *Graph Theory and Applications, Lecture Notes in Math.*, Vol. 303 (Springer, New York, 1972) 99–110.
- [12] R. Graham and P. Winkler, On isometric embeddings of graphs, *Trans. Amer. Math. Soc.* 288 (1985) 527–536.
- [13] W. Imrich, Embedding graphs into Cartesian products, *Ann. New York Acad. Sci.* 576 (1989) 266–274.
- [14] E. Wilkeit, Isometric embeddings in Hamming graphs, *J. Combin. Theory Ser. B* 50 (1990) 179–197.
- [15] P. Winkler, Proof of the squashed cube conjecture, *Combinatorica* 3 (1983) 135–139.
- [16] P. Winkler, Isometric embeddings in products of complete graphs, *Discrete Appl. Math.* 7 (1984) 221–225.
- [17] P. Winkler, The metric structure of a graph, in: C. Whitehead, ed., *Theory and Applications, London Math. Soc. Lecture Note Ser.*, Vol. 123 (Cambridge University Press, Cambridge, 1987) 197–221.