

Recognizing halved cubes in a constant time per edge

Wilfried Imrich

Department of Mathematics and Applied Geometry
Montanuniversität Leoben
A-8700 Leoben, Austria

Sandi Klavžar* and Aleksander Vesel
Department of Mathematics, PEF
University of Maribor
Koroška cesta 160
62000 Maribor, Slovenia

*This work was supported in part by the Ministry of Science and Technology of Slovenia under the grant P1-0206-101.

Proposed running head: Recognizing halved cubes

Mailing address:

Sandi Klavžar
Department of Mathematics, PEF
University of Maribor
Koroška cesta 160
62000 Maribor, Slovenia
e-mail: sandi.klavzar@uni-lj.ac.mail.si

Abstract

Graphs that can be isometrically embedded into the metric space ℓ_1 are called ℓ_1 -graphs. Halved cubes play an important role in the characterization of ℓ_1 -graphs. We present an algorithm that recognizes halved cubes in $O(n \log^2 n)$ time.

1 Introduction

Hamming graphs are exactly those graphs which can be isometrically embedded into a Cartesian product of complete graphs. In case every one of the factors is the complete graph K_2 on two vertices one obtains an isometric embedding into a hypercube and speaks of a binary Hamming graph.

If one relaxes the condition of isometry and considers so-called scale embeddings into hypercubes a class larger than that of Hamming graphs arises. It has been characterized by Assouad and Deza [1] as the class of graphs isometrically embeddable into the metric space ℓ_1 .

By a scale embedding of a graph G into a graph H we mean a mapping

$$\psi : V(G) \rightarrow V(H)$$

for which there exists a positive integer λ such that

$$d_H(\psi(u), \psi(v)) = \lambda d_G(u, v)$$

for all $u, v \in V(G)$, where d_H and d_G denote the usual path distance in G and H , respectively. It has been proved by Deza and Grishukhin [4] and Shpectorov [13] that a graph G is an ℓ_1 -graph if and only if it is an isometric subgraph of a Cartesian product of complete graphs, cocktail party graphs or halved cubes and that ℓ_1 -graphs can be recognized in polynomial time.

Aurenhammer, Formann, Idury, Schäffer and Wagner [2] (see also Imrich and Klavžar [11]) proved that it can be decided in $O(mn)$ time whether a given graph on n vertices and m edges is Hamming graph. Thus, the question whether this is also true for ℓ_1 -graphs arises. We cannot answer this question, in particular we cannot efficiently decide whether a graph is an isometric subgraph of a halved cube. However, as a first step, we are able to recognize halved cubes in $O(n \log^2 n)$ steps, where n denotes the number of vertices.

2 Recognizing halved cubes

Let Q_d be the d -cube and let $V_1 \cup V_2$ be its bipartition as a bipartite graph. Then the *halved cube* Q'_d is the graph with $V(Q'_d) = V_1$, where u is adjacent to v in Q'_d if $d_{Q_d}(u, v) = 2$. Clearly, Q'_d has 2^{d-1} vertices and is $\binom{d}{2}$ regular

TWO-CLIQUEs (uv, Q, Q');

1. let $Y = N(u) \cap N(v)$; if $|Y| \neq 2(d-2)$ then reject;
2. $Q := \{u, v\}$; $Q' := \{u, v\}$;
3. let w_1, w_2 and w_3 be any three vertices from Y and let G_1 be the subgraph induced by w_1, w_2 and w_3 .
 - 3.1 if G_1 has no edge then reject;
 - 3.2 if G_1 has three edges then $Q := Q \cup \{w_1, w_2, w_3\}$, $Y := Y - \{w_1, w_2, w_3\}$;
 - 3.3 if G_1 has one edge, say w_1w_2 , then $Q := Q \cup \{w_1, w_2\}$, $Y := Y - \{w_1, w_2\}$;
 - 3.4 if G_1 is a path on three vertices, say $w_1w_2w_3$, then let w_4 be an additional vertex of Y and let G_2 be the subgraph induced by w_1, w_2, w_3 and w_4 .
 - 3.4.1 if G_2 has three edges and w_4 is adjacent to either w_1 or w_3 , say w_3 , then $Q := Q \cup \{w_1, w_2\}$, $Q' := Q' \cup \{w_3, w_4\}$, $Y := Y - \{w_1, w_2, w_3, w_4\}$;
 - 3.4.2 if G_2 has four edges and w_4 is adjacent to either w_1 and w_2 or to w_2 and w_3 , say to w_1 and w_2 , then $Q := Q \cup \{w_1, w_2, w_4\}$, $Q' := Q' \cup \{w_3\}$, $Y := Y - \{w_1, w_2, w_3, w_4\}$;
 - 3.4.3 if G_2 is the 4-cycle C_4 then let w_5 be an additional vertex of Y ;
 - 3.4.3.1 if w_5 is adjacent to exactly two adjacent vertices of G_2 , say w_1 and w_2 , then $Q := Q \cup \{w_1, w_2, w_5\}$, $Q' := Q' \cup \{w_3, w_4\}$, $Y := Y - \{w_1, w_2, w_3, w_4, w_5\}$;
 - 3.4.3.2 in all the other cases reject;
 - 3.4.4 if none of the above cases occur then reject;
4. for any vertex $w \in Y$ do: if w is adjacent to both w_1 and w_2 then $Q := Q \cup \{w\}$ else $Q' := Q' \cup \{w\}$;
5. if $|Q| \neq d$ then reject;
6. if $|E(Q)| < \binom{d}{2}$ or $|E(Q')| < \binom{d}{2}$ then reject.

We next describe a procedure which computes all d -cliques of G . The cliques of a graph will be consecutively indexed and for an edge uv let the variable $l_{u,v}$ be defined by

$$l_{u,v} = \begin{cases} 0; & uv \text{ does not belong to any clique yet,} \\ i; & uv \text{ belongs to a clique indexed } i, \\ -1; & uv \text{ belongs to two cliques.} \end{cases}$$

CLIQUEES (Q_1, Q_2, \dots, Q_n);

1. If G is not a connected, $\binom{d}{2}$ -regular graph on 2^{d-1} vertices for some $d \geq 5$, then reject.
2. $i := 0$; {clique counter}
3. for all edges $uv \in E(G)$ do $l_{u,v} := 0$;
4. while there is an edge $uv \in E(G)$ with $l_{u,v} \neq -1$ and $i < n = 2^{d-1}$ do:
 - 4.1. TWO-CLIQUEES (uv, Q, Q');
 - 4.2. if $l_{u,v} = 0$ then
 - $Q_{i+1} := Q; Q_{i+2} := Q'$;
 - for all $wz \in E(Q_{i+1})$ do
 - if $l_{w,z} = 0$ then $l_{w,z} := i + 1$ else
 - if $l_{w,z} > 0$ then $l_{w,z} := -1$, otherwise reject;
 - for all $wz \in E(Q_{i+2})$ do
 - if $l_{w,z} = 0$ then $l_{w,z} := i + 2$ else
 - if $l_{w,z} > 0$ then $l_{w,z} := -1$, otherwise reject;
 - $i := i + 2$
 - else
 - if $Q = Q_{l_{u,v}}$ then $Q_{i+1} := Q'$ else
 - if $Q' = Q_{l_{u,v}}$ then $Q_{i+1} := Q$ else reject;
 - for all $wz \in E(Q_{i+1})$ do
 - if $l_{w,z} = 0$ then $l_{w,z} := i + 1$ else
 - if $l_{w,z} > 0$ then $l_{w,z} := -1$, otherwise reject;
 - $i := i + 1$;
5. If $i < n$ then reject.

We have arrived to the procedure for recognizing halved cubes.

HALVED-CUBE (G);

1. CLIQUES (Q_1, Q_2, \dots, Q_n);
2. Let H be a graph which we get from G in the following way. To every d -clique Q of G we add a new vertex and join it to every vertex of Q . These are the newly defined edges of H . The original edges of G are all removed.
3. If H is Q_d then G is Q'_d , otherwise reject.

Theorem 1 *Let G be a graph on n vertices. Then it can be decided in $O(n \log^2 n)$ time whether G is a halved cube.*

Proof. The correctness of the algorithm follows from the previous discussion.

As G is $\binom{d}{2}$ -regular, Step 1 of TWO-CLIQUES can be performed in $O(d^2)$ time if the input is a preordered adjacency list. Steps 2, 3 and 5 can be computed in a constant time and Step 4 needs $O(d)$ time. Finally, Step 6 can be performed in $O(d^2)$ time. Hence the total complexity of TWO-CLIQUES is $O(d^2) = O(\log^2 n)$.

Whenever we perform Step 4 of CLIQUES, at least one new d -clique is added to the list of cliques. Thus Step 4 is executed at most 2^{d-1} times. Since each call of TWO-CLIQUES requires $O(d^2)$ time and Step 4.2. can clearly be performed within the same time, the time complexity of CLIQUES is $O(n \log^2 n)$.

Finally, hypercubes can be recognized in $O(n \log n)$ time using an algorithm from [3]. □

Another approach to recognize hypercubes efficiently is the following. In [12] Jha and Slutzky gave an algorithm which recognizes median graphs in $O(n^2 \log n)$ time. As it was observed by Hagauer [8], a slight modification of this algorithm can be used to recognize hypercubes in $O(n \log n)$ time. We will not give the details here, but wish to mention that the improvement in the complexity is due to the observation that we don't need to take care of convexity (which is the bottleneck in recognizing median graphs, cf. also [9]).

Concluding remark

In fact, without going into details, we should like to mention that by a result of Graham and Winkler [7] about so-called canonical isometric embeddings of graphs into Cartesian products together with an algorithm of Feder [6], a good algorithm for recognizing isometric subgraphs of halved cubes would suffice for a good algorithm for recognizing ℓ_1 -graphs. Unfortunately such an algorithm is not yet known, cf. Deza and Laurent [5].

References

- [1] P. Assouad and M. Deza, *Metric subspaces of L^1* , Publ. Math. Orsay, vol. 3, 1982.
- [2] F. Aurenhammer, M. Formann, R. Idury, A. Schäffer and F. Wagner, Faster isometric embeddings in products of complete graphs, *Discrete Appl. Math.*, to appear.
- [3] K.V.S. Bhat, On the complexity of testing a graph for n -cube, *Inform. Process. Lett.* **11** (1980) 16–19.
- [4] M. Deza and V.P. Grishukhin, Hypermetric graphs, *Quart. J. Math. Oxford Ser. (2)* **44** (1993), 399–433.
- [5] M. Deza and M. Laurent, ℓ_1 -rigid graphs, *J. Algebraic Combin.* **3** (1994), 153–175.
- [6] T. Feder, Product graph representations, *J. Graph Theory* **16** (1992), 467–488.
- [7] R. L. Graham and P. M. Winkler, On isometric embeddings of graphs, *Trans. Amer. Math. Soc.* **288** (1985), 527–536.
- [8] J. Hagauer, personal communication.
- [9] J. Hagauer, W. Imrich and S. Klavžar, Recognizing graphs of windex 2, Preprint Series Univ. Ljubljana **31** (1993) 410.
- [10] J. Hemmeter, Distance-regular graphs and halved graphs, *European J. Combin.* **7** (1986), 119–129.
- [11] W. Imrich and S. Klavžar, A simple $O(mn)$ algorithm for recognizing Hamming graphs, *Bull. Inst. Comb. Appl.* **9** (1993), 45–56.
- [12] P.K. Jha and G. Slutzki, Convex-expansion algorithms for recognition and isometric embedding of median graphs, *Ars Combin.* **34** (1992), 75–92.
- [13] S.V. Shpectorov, On scale embeddings of graphs into hypercubes, *European J. Combin.* **14** (1993), 117–130.